

## Tutorial-2, Statistical Mechanics & Others (Paper-203)

Dr. R K Hazra

Maximum Marks: 50

Q-1. Obtain partition function  $Z(N, V, E)$  of a microcanonical ensemble.

(Hint: use multinomial expansion with degeneracy factor)

Q-2. Energy of particle in a cube of side ' $a$ ',  $\epsilon_{n_x, n_y, n_z} = \frac{\pi^2 \hbar^2}{2ma^2} (n_x^2 + n_y^2 + n_z^2)$ . Find approximate total degeneracies of states with very high energy.

Q-3. Calculate number of states ( $\omega$ ) between  $\epsilon$  and  $\epsilon + \Delta\epsilon$  for particle in a cube with very high energy. Find  $\omega$  for  $T = 300K$ ,  $m = 10^{-22}g$ ,  $a = 10cm$  and  $\Delta\epsilon = 0.01\epsilon$

Q-4. Given  $dE = TdS - PdV$  for a one component reversible system, find relations of  $T$  and  $P$  with  $E$  respectively. Obtain further relation between  $T$  and  $P$ .

Q-5. Given  $dH = TdS + VdP$  for a one component reversible system, find relations of  $T$  and  $V$  with  $H$  respectively. Obtain further relation between  $T$  and  $V$ .

Q-6. Given  $dG = VdP - SdT$  for a one component reversible system, find relations of  $V$  and  $S$  with  $G$  respectively. Obtain further relation between  $V$  and  $S$ .

Q-7. Given  $dA = -PdV - SdT$  for a one component reversible system, find relations of  $P$  and  $S$  with  $A$  respectively. Obtain further relation between  $P$  and  $S$ .

Q-8. Define  $E$ ,  $H$ ,  $A$  and  $G$  for many-component closed system. Illustrate chemical potential ( $\mu$ ) in terms of above functions.

Q-9. Using method of Lagrangian Multipliers, obtain maximum ground state level of a particle in 3-D box with sides  $a$ ,  $b$  and  $c$  having total volume constant.

Q-10. Using method of Lagrangian Multipliers, obtain maximum perimeter of a rectangle having constant area.

Books: McQuarrie (Statistical Mechanics), Callen (Thermodynamics and Thermostatistics), Nash (Elements of Statistical Thermodynamics), Atkins (Physical Chemistry), Landau & Lifshitz (Statistical Physics).